

1. A particle moves three times, counterclockwise, along the circle

$$x^2 + (y - 1)^2 = 9.$$

If the particle starts at the point $(3, 1)$, then the parametric equations of the path of the particle are

- (a) $x = 3 \cos t, y = 1 + 3 \sin t, 0 \leq t \leq 6\pi$
 (b) $x = 3 \cos t, y = 1 - 3 \sin t, 0 \leq t \leq 6\pi$
 (c) $x = 3 \sin t, y = 1 + 3 \cos t, 0 \leq t \leq 6\pi$
 (d) $x = 3 \cos(3t), y = 1 - 3 \sin(3t), 0 \leq t \leq 2\pi$
 (e) $x = 3 \sin(3t), y = 1 + 3 \cos(3t), 0 \leq t \leq 2\pi$

$$x = 3 \cos t, y = 1 + 3 \sin t$$

$$0 \leq t \leq 2\pi$$

$$\text{At } t=0, (x, y) = (3, 1)$$

As t increases from 0 to 6π , particle moves 3 times counterclockwise.

10.1
33(b)

2. The curve given parametrically by

$$x = \tan \theta, y = 2 \sec \theta, -\pi/2 < \theta < \pi/2$$

is

- (a) a hyperbola
 (b) an ellipse
 (c) a parabola with horizontal axis of symmetry
 (d) a parabola with vertical axis of symmetry
 (e) a straight line

$$x^2 = \frac{y^2}{4} - 1$$

10.1
Similar 18

3. The curve given by the parametric equations

$$x = 2 \cos^3 \theta,$$

$$y = 2 \sin^3 \theta, \quad 0 \leq \theta < \frac{7\pi}{4}$$

has

$$\frac{dx}{d\theta} = -6 \cos^2 \theta \sin \theta$$

$$\frac{dy}{d\theta} = 6 \sin^2 \theta \cos \theta$$

$$\frac{dy}{dx} = -\tan \theta = 0$$

$$\Rightarrow \theta = 0, \pi$$

- (a) only one horizontal and only one vertical tangent lines
- (b) only one horizontal tangent line and no vertical tangent line
- (c) no horizontal tangent line and only one vertical tangent line
- (d) two horizontal and one vertical tangent lines
- (e) one horizontal and two vertical tangent lines

10.2
Similar 20(b)

$$\theta = 0, \pi, y = 0$$

$$\theta = \frac{\pi}{2}, \frac{3\pi}{2}, x = 0$$

$$\frac{dy}{dx} = \pm \infty \Rightarrow \theta = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$\left(\frac{dy}{dx} \rightarrow \pm \infty \text{ as } \theta \rightarrow \frac{\pi}{2}^+, \frac{\pi}{2}^-, \frac{3\pi}{2}^+, \frac{3\pi}{2}^-\right)$$

A* $\theta = 0$ as well as $\theta = \pi \Rightarrow$ one horizontal line ($y=0$)
 At $\theta = \frac{\pi}{2}, \frac{3\pi}{2} \Rightarrow$ one vertical tangent line ($x=0$)

4. The curve given by the parametric equations

$$x = t^3 - 3t \text{ and } y = t^2$$

is

- (a) concave up when $-1 < t < 1$
- (b) concave down when $t < 0$
- (c) concave up when $t > 0$
- (d) concave down when $-\frac{1}{2} < t < \frac{1}{2}$
- (e) concave up when $t > 1$

$$\frac{dy}{dx} = \frac{2t}{3(t^2-1)} = 0$$

$$\frac{dy}{dx} = 0 \Rightarrow t = 0, \quad \frac{dx}{dt} = 0 \Rightarrow t = \pm 1$$

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt}\left(\frac{dy}{dx}\right)}{\frac{dx}{dt}} = \frac{6(t^2-1) - (2t)^2}{9(t^2-1)^2}$$

$$= \frac{-2(t^2+1)}{9(t^2-1)^3}$$

$$\frac{d^2y}{dx^2} > 0, \text{ when } -1 < t < 1 \quad \text{CU}$$

$$< 0, \text{ when } t > 1 \text{ or } t < -1 \quad \text{CD}$$

10.2
Example P650

5. The area of the region enclosed by the curve that is given by the parametric equations

$$x = 2 \cos \theta, y = \frac{1}{2} \sin \theta, 0 \leq \theta \leq 2\pi, \text{ is}$$

- (a) π
 (b) 3π
 (c) 0
 (d) 2π
 (e) 4π

$$\begin{aligned} A &= 4 \int_0^2 y dx && x=0 \Rightarrow \theta = \pi/2 \\ &= 4 \int_0^{\pi/2} \frac{1}{2} \sin \theta (-2 \sin \theta) d\theta && x=2 \Rightarrow \theta = 0 \\ &= 4 \int_0^{\pi/2} \sin^2 \theta d\theta = \int_0^{\pi/2} (2 - 2 \cos 2\theta) d\theta \\ &= (2\theta - \sin 2\theta) \Big|_0^{\pi/2} \\ &= (\pi - 0) - (0 - 0) = \pi \end{aligned}$$

$$\frac{10.2}{\text{a. } 31}$$

6. The surface area of the solid generated by rotating the curve given by the parametric equations

$$x = 6t, y = 3t^2, 0 \leq t \leq 1,$$

about the y -axis is

- (a) $24\pi [2\sqrt{2} - 1]$
 (b) $12\pi [\sqrt{2} - 1]$
 (c) $9\pi [1 - 2\sqrt{2}]$
 (d) $6\pi [\sqrt{2} + 1]$
 (e) $3\pi [2 - \sqrt{2}]$

$$\begin{aligned} ds &= \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \sqrt{36 + 36t^2} dt \\ &= 6\sqrt{1+t^2} dt \\ S &= \int_0^1 2\pi x ds \\ &= \int_0^1 2\pi \cdot 6t \cdot 6\sqrt{1+t^2} dt \\ &= 36\pi \int_0^1 2t\sqrt{1+t^2} dt \\ &= 36\pi \int_0^1 u^{1/2} du && 1+t^2 = u \\ &= 36\pi \left[\frac{2}{3} (1+t^2)^{3/2} \right]_0^1 && 2t dt = du \\ &= 24\pi [2^{3/2} - 1] \\ &= 24\pi [2\sqrt{2} - 1] \end{aligned}$$

$$\frac{10.2}{\text{a. } 65}$$

7. The polar equation $r = \tan \theta \sec \theta$ represents a

- (a) Parabola
- (b) Circle
- (c) Line
- (d) Ellipse
- (e) Hyperbola

$$r = \frac{\sin \theta}{\cos \theta} \cdot \frac{1}{\cos \theta}$$

$$r \cos^2 \theta = \sin \theta$$

$$r^2 \cos^2 \theta = r \sin \theta$$

$$x^2 = y$$

$$\Rightarrow \text{Parabola}$$

10.3
Q.18

8. The graph of the polar curve $r = 3 \cos \theta$, has a vertical tangent line when $(r, \theta) =$

- (a) $(0, \pi/2)$
- (b) $(0, \pi)$
- (c) $(0, \pi/4)$
- (d) $(3, \pi/2)$
- (e) $(0, 2\pi)$

$$x = r \cos \theta$$

$$= 3 \cos^2 \theta$$

$$\frac{dx}{d\theta} = -6 \cos \theta \sin \theta = 0$$

$$\Rightarrow \sin 2\theta = 0$$

$$\Rightarrow \theta = 0, \frac{\pi}{2}$$

$$\text{At } \theta = 0, r = 3$$

$$\text{At } \theta = \pi/2, r = 0$$

$$(r, \theta) = (0, \frac{\pi}{2})$$

10.3
Q.61