

Q1.

A standing wave having three nodes is set up in a string fixed at both ends. If the frequency of the wave is doubled, how many antinodes will there be?

- A) 4
- B) 3
- C) 2
- D) 5
- E) 6

---

Q2.

A stationary motion detector sends sound waves of frequency 0.120 MHz towards a truck approaching (the detector) at a speed of 50.0 m/s. What is the frequency of the waves reflected back to the detector? [Speed of sound = 343 m/s]

- A) 0.161 MHz
- B) 0.103 MHz
- C) 0.140 MHz
- D) 0.234 MHz
- E) 0.186 MHz

---

Q3.

A box with a total surface area of  $1.20 \text{ m}^2$  and a wall thickness of 4.00 cm is made of an insulating material. A 10.0 W electric heater inside the box maintains the inside temperature at  $15.0 \text{ }^\circ\text{C}$  above the outside temperature. Find the thermal conductivity of the insulating material.

- A) 0.022 W/m.K
- B) 2.20 W/m.K
- C) 0.034 W/m.K
- D) 0.016 W/m.K
- E) 1.23 W/m.K

---

Q4.

An ideal gas initially at 330 K is compressed at a constant pressure of  $25.0 \text{ N/m}^2$  from a volume of  $3.0 \text{ m}^3$  to a volume of  $1.00 \text{ m}^3$ . In the process, 75.0 J is lost by the gas as heat. What is the change in internal energy of the gas?

- A)  $-25.0 \text{ J}$
- B)  $-125 \text{ J}$
- C)  $+50.0 \text{ J}$
- D)  $+65.0 \text{ J}$
- E)  $-75.0 \text{ J}$

---

Q5.

A Carnot engine has an efficiency of  $\epsilon$  when operating between  $T_H = 400 \text{ }^\circ\text{C}$  and  $T_L = 200 \text{ }^\circ\text{C}$ . What will be the efficiency of the same Carnot engine when operating between  $T_H = 800 \text{ }^\circ\text{C}$  and  $T_L = 400 \text{ }^\circ\text{C}$ .

- A)  $1.25\epsilon$

- B)  $2.0\epsilon$
- C)  $0.50\epsilon$
- D)  $1.5\epsilon$
- E)  $0.77\epsilon$

Q6.

Two identical 0.20 kg masses are placed 1.0 m apart (center to center) on a frictionless surface. Each has  $+10 \mu\text{C}$  of charge. What is the initial acceleration of one of the masses if it is released from rest and allowed to move?

- A)  $4.5 \text{ m/s}^2$
- B)  $2.3 \text{ m/s}^2$
- C)  $6.3 \text{ m/s}^2$
- D)  $1.4 \text{ m/s}^2$
- E)  $5.2 \text{ m/s}^2$

Q7.


Two point charges  $q_1 = +15 \mu\text{C}$  and  $q_2 = -10 \mu\text{C}$  are placed in  $xy$  plane. If  $q_1$  is placed at (20 cm, 0) and  $q_2$  is placed at (0, 10 cm), find the resultant electric field at the origin.

- A)  $(-3.38 \times 10^6 \hat{i} + 9.00 \times 10^6 \hat{j}) \text{ N/C}$
- B)  $(3.38 \times 10^6 \hat{i} + 9.00 \times 10^6 \hat{j}) \text{ N/C}$
- C)  $(-3.38 \times 10^6 \hat{i} - 9.00 \times 10^6 \hat{j}) \text{ N/C}$
- D)  $(-9.00 \times 10^6 \hat{i} + 3.38 \times 10^6 \hat{j}) \text{ N/C}$
- E)  $(9.00 \times 10^6 \hat{i} + 3.38 \times 10^6 \hat{j}) \text{ N/C}$

Q8.

**FIGURE 1** shows portions of a large non-conducting sheet placed in parallel with long line of charge having a uniform charge per unit length  $\lambda$ . The surface charge density of the non-conducting sheet is  $\sigma = +4.0 \mu\text{C}/\text{m}^2$ . If the electric field intensity at point P, 2.0 cm on right of the line charge, is zero, find the linear charge density  $\lambda$ .

Fig#



- A)  $-0.25 \mu\text{C}/\text{m}$
- B)  $+0.25 \mu\text{C}/\text{m}$
- C)  $+0.38 \mu\text{C}/\text{m}$
- D)  $-0.38 \mu\text{C}/\text{m}$
- E)  $-0.63 \mu\text{C}/\text{m}$

Q9.

A thick spherical conducting shell of outer radius 5.0 cm has a net charge  $Q = +10 \mu\text{C}$ . A point charge of  $-3.0 \mu\text{C}$  is placed at its center. Find the surface charge density on the **outer surface** of the shell.

- A)  $223 \mu\text{C}/\text{m}^2$
- B)  $414 \mu\text{C}/\text{m}^2$
- C)  $318 \mu\text{C}/\text{m}^2$
- D)  $203 \mu\text{C}/\text{m}^2$
- E)  $196 \mu\text{C}/\text{m}^2$

Q10.


Two conducting spherical shells of same outer radius  $r = 5.0 \text{ cm}$  are placed at center to center distance of 20 cm. The charge of  $+10 \mu\text{C}$  and  $-20 \mu\text{C}$  is uniformly distributed over the outer surface of sphere 1 and 2, respectively, find the total potential at the center of sphere 1. (Assume potential is zero at infinity)

- A) +900 kV
- B) +270 kV
- C) -900 kV
- D) +450 kV
- E) +600 kV

Q11.

In **FIGURE 2**  $C_1 = 2.0 \mu\text{F}$ ,  $C_2 = 4.0 \mu\text{F}$ , and  $C_3 = 6.0 \mu\text{F}$ . Find the equivalent capacitance between points a and b.

Fig#



- A)  $8.7 \mu\text{F}$
- B)  $18 \mu\text{F}$
- C)  $9.8 \mu\text{F}$
- D)  $12 \mu\text{F}$
- E)  $2.0 \mu\text{F}$

Q12.


A current of 3.20 A exist in a copper wire whose diameter is 4.00 mm. The number of charge carrier per unit volume is  $8.49 \times 10^{28} \text{ m}^{-3}$ . Assuming the current density is uniform, calculate the electron drift speed.

- A)  $1.87 \times 10^{-5} \text{ m/s}$
- B)  $7.49 \times 10^{-5} \text{ m/s}$
- C)  $4.25 \times 10^{-6} \text{ m/s}$
- D)  $1.40 \times 10^{-5} \text{ m/s}$
- E)  $3.23 \times 10^{-5} \text{ m/s}$

Q13.

Two real batteries with some internal resistances are shown in **FIGURE 3**. Find the potential difference between points A and B ( $V_A - V_B$ ).

Fig#




- A) 13.0 V
- B) 23.5 V
- C) 11.8 V
- D) 24.5 V
- E) 18.0 V

Q14.

In a circuit shown in **FIGURE 4** if the potential of point **A** is zero find the potential of point **B**.

Fig#




- A) -2.0 V
- B) +2.0 V
- C) +6.0 V
- D) -6.0 V
- E) -4.0 V

Q15.

For the circuit given in **FIGURE 5**, if the current through one of the  $6.00 \Omega$  is  $0.500 \text{ A}$  find the current through the  $4.00 \Omega$  resistor.

Fig#




- A) 2.25 A
- B) 0.33 A
- C) 3.50 A
- D) 1.55 A
- E) 0.50 A

Q16.

If the current through the  $150\ \Omega$  resistor is  $1.00\ \text{A}$  as shown in **FIGURE 6**, find the current through the  $51.0\ \Omega$  resistor.

Fig#




- A)  $8.47\ \text{A}$
- B)  $3.00\ \text{A}$
- C)  $5.42\ \text{A}$
- D)  $11.4\ \text{A}$
- E)  $6.45\ \text{A}$

Q17.

Switch  $S$  in **FIGURE 7** is closed at time  $t = 0$ , to begin charging an initially uncharged capacitor  $C = 15.0\ \mu\text{F}$  through a resistor  $R = 20.0\ \Omega$ . At what time is the potential difference across the capacitor double to that across the resistor?

Fig#




- A)  $3.30 \times 10^{-4}\ \text{s}$
- B)  $2.08 \times 10^{-4}\ \text{s}$
- C)  $1.25 \times 10^{-4}\ \text{s}$
- D)  $1.04 \times 10^{-4}\ \text{s}$
- E)  $0.52 \times 10^{-4}\ \text{s}$

Q18.

A proton enters with the same speed in four regions of uniform magnetic fields of same magnitudes but in different directions, as shown in **FIGURE 8**. Rank regions according to magnitude of the force on the protons, **GREATEST FIRST**.

Fig#




- A) a and b tie, d, c
- B) b and c tie, a, d
- C) b and c tie, d, a
- D) a, then b and c tie, d
- E) a, d, b, c

Q19.

In **FIGURE 9** a charged particle electron or proton (you must decide which) moves in uniform magnetic field  $\vec{B}$  (out of the page), goes through half circle, and then exits that region. The charged particle spends 240 ns in the region, find the magnitude of magnetic field.

Fig#



- A) 0.137 T
- B) 0.298 T
- C)  $1.49 \times 10^{-4}$  T
- D)  $3.72 \times 10^{-4}$  T
- E) 0.579 T

Q20.

An 80.0 cm long wire, laying along the positive x axis (with one end at the origin), carries a current of 0.80 A in the negative x direction and placed in a magnetic field.  $\vec{B} = 4.0\hat{i} + 12\hat{j}$ , where x in meters and B in mT. Find, in unit vector notation, the magnetic force on the wire?


- A)  $(-7.68 \times 10^{-3} \hat{k})N$
- B)  $(+2.62 \times 10^{-3} \hat{k})N$
- C)  $(-2.62 \times 10^{-3} \hat{k})N$
- D)  $(+5.62 \times 10^{-3} \hat{j})N$

E)  $(-4.72 \times 10^{-3} \hat{i})N$

Q21.

A rectangular loop with height  $h = 6.50$  cm and width  $w = 5.40$  cm is in a uniform magnetic field of magnitude  $B = 0.250$  T, which points in negative  $y$  direction as shown in **FIGURE 10**. The loop makes an angle of  $\theta = 30^\circ$  with the  $y$  axis and carries a current of  $9.00$  A in the direction indicated. What is the magnitude of the torque on the loop?

Fig#




- A)  $6.84 \times 10^{-3}$  N.m
- B)  $9.56 \times 10^{-3}$  N.m
- C)  $7.62 \times 10^{-3}$  N.m
- D)  $3.95 \times 10^{-3}$  N.m
- E)  $4.32 \times 10^{-3}$  N.m

Q22.

Two concentric circular loops of radii  $r_1 = 20.0$  cm and  $r_2 = 30.0$  cm are located in  $xy$  plane; each carries current of same magnitude  $i = 7.00$  A but in opposite direction as shown in **FIGURE 11**. Find the net magnetic dipole moment of the loops.


Fig#



- A)  $1.10 \text{ A}\cdot\text{m}^2$  into the page
- B)  $2.86 \text{ A}\cdot\text{m}^2$  into the page
- C)  $1.10 \text{ A}\cdot\text{m}^2$  out of the page
- D)  $2.86 \text{ A}\cdot\text{m}^2$  out of the page
- E)  $1.98 \text{ A}\cdot\text{m}^2$  into the page

Q23.

Wire 1 and wire 2 placed parallel to y axis at  $x = 0$  and  $x = 10 \text{ cm}$ , respectively, and carries currents  $i_1 = 4.0\text{A}$  and  $i_2 = 8.0\text{A}$  in opposite directions as shown in **FIGURE 12**. A third wire carries current in positive y direction is to be placed parallel to wire 1 and wire 2 such that net force per unit length on wire 3 due to wire 1 and wire 2 is zero. Find the position  $x$  of wire 3.




- A)  $-10 \text{ cm}$
- B)  $+20 \text{ cm}$
- C)  $+15 \text{ cm}$
- D)  $-5.0 \text{ cm}$
- E)  $-15 \text{ cm}$

Q24.

In **FIGURE 13** wire 1 consist of a circular arc of radius  $R$  with central angle of  $\theta = 120^\circ$  and two radial lengths and carries current  $i_1 = 2.00\text{A}$  in the direction indicated. Wire 2 is long and straight and it carries a current  $i_2$  and placed at a distance of  $R/2$  from the center of circular arc. If the net magnetic field at the center of the arc  $O$  is zero find the current  $i_2$ .

Fig#



- A) 1.05 A
- B) 2.75 A
- C) 0.84 A
- D) 1.34 A
- E) 2.05 A


Q25.

**FIGURE 14** shows a cross section across the diameter of two long cylindrical conducting wires 1 and 2 of same radius  $R$  carrying same uniform current but in opposite directions. Four square paths (of side length  $R$ ) of same dimensions are indicated for the line integral  $\oint \vec{B} \cdot d\vec{s}$ .

Rank the paths according to the magnitude of  $\oint \vec{B} \cdot d\vec{s}$  taken in the directions shown,

**GREATEST FIRST.**

Fig#




- A) 4, then 1 and 3 tie, 2
- B) 1 and 3 tie, 2, 4
- C) 4, then 1, 2, and 3 tie
- D) 4, 3, 2, 1
- E) 3, 4, 1, 2

Q26.

A circular wire loop has radius of 0.12 m and carries current  $i = 0.10\text{A}$  placed in the  $xy$  plane in a uniform magnetic field  $\vec{B} = -1.5\hat{k}\text{T}$ , as shown in **FIGURE 15**. Find the potential energy of the loop in the position shown.

Fig#



- A)  $+6.79 \times 10^{-3}\text{ J}$
- B)  $-6.79 \times 10^{-3}\text{ J}$
- C)  $+5.65 \times 10^{-2}\text{ J}$
- D) 0
- E)  $-5.65 \times 10^{-2}\text{ J}$

Q27.


A solenoid has a length  $L = 1.55\text{ m}$  and an inner diameter  $d = 4.15\text{ cm}$ , and carries a current  $i = 4.80\text{ A}$ . The solenoid consists of six close-packed layers, each with 750 turns along length  $L$ . What is the magnitude of magnetic field at its center?

- A) 17.5 mT
- B) 23.8 mT
- C) 5.65 mT
- D) 13.5 mT
- E) 19.8 mT

Q28.

**FIGURE 16** shows three situations in which a wire loop lies partially in a uniform magnetic field. The magnetic field is either increasing or decreasing, as indicated. In each situation, a battery is part of the loop. In which situation(s) is/are the induced emf and the battery emf in the same direction along the loop.

Fig#



- A) b only  
B) b and c  
C) a, b, and c  
D) c only  
E) a and c

Q29.

The magnetic flux through a loop increases according to the relation  $\Phi_B = 6.0t^2 + 7.0t$ , where  $\Phi_B$  is in  $\text{mT}\cdot\text{m}^2$  and  $t$  is in seconds. What is the magnitude of the emf induced in the loop when  $t = 2.0$  s?

- A) 31 mV  
B) 38 mV  
C) 24 mV  
D) 17 mV  
E) 40 mV

Q30.

A 60.0 cm copper wire is formed into a square loop and placed perpendicular to a uniform magnetic field that is increasing at the constant rate 12.0 mT/s. If the resistance of the loop is 20.0  $\Omega$ , at what rate is thermal energy generated in the loop?

- A)  $3.65 \times 10^{-9}$  W  
B)  $2.70 \times 10^{-9}$  W  
C)  $7.45 \times 10^{-9}$  W  
D)  $1.35 \times 10^{-9}$  W  
E)  $4.12 \times 10^{-9}$  W