

Q1.

For the wave described by  $y = 0.02 \sin(kx)$  at  $t = 0$  s, the first maximum displacement at a positive  $x$  coordinate occurs at  $x = 4$  m. Where on the positive  $x$  axis does the second maximum displacement occur? ( $x$  and  $y$  are measured in m)

- A) 20 m
- B) 12 m
- C) 34 m
- D) 48 m
- E) 59 m

Q2.


The sound intensity 5.00 m from a point source is  $0.500 \text{ W/m}^2$ . The power output of the source is:

- A) 157 W
- B) 391 W
- C) 710 W
- D) 235 W
- E) 458 W

Q3.

**Figure 1** shows four vertical thermometers, labeled W, X, Y, and Z. The freezing and boiling points of water are indicated. Rank the 65 degree temperatures on different scale from **highest to lowest**.

Fig#



- A)  $65^\circ\text{Z}$ ,  $65^\circ\text{W}$ ,  $65^\circ\text{X}$ ,  $65^\circ\text{Y}$
- B)  $65^\circ\text{Z}$ ,  $65^\circ\text{X}$ ,  $65^\circ\text{W}$ ,  $65^\circ\text{Y}$
- C)  $65^\circ\text{W}$ ,  $65^\circ\text{X}$ ,  $65^\circ\text{Y}$ ,  $65^\circ\text{Z}$
- D)  $65^\circ\text{Y}$ ,  $65^\circ\text{W}$ ,  $65^\circ\text{X}$ ,  $65^\circ\text{Z}$
- E)  $65^\circ\text{X}$ ,  $65^\circ\text{Y}$ ,  $65^\circ\text{Z}$ ,  $65^\circ\text{W}$

Q4.

The approximate number of air molecules in a  $1.00 \text{ m}^3$  volume at room temperature (300 K) and atmospheric pressure is: (Assume air to be an ideal gas)

- A)  $2.44 \times 10^{25}$
- B)  $3.30 \times 10^{25}$
- C)  $1.63 \times 10^{25}$

- D)  $4.71 \times 10^{25}$   
E)  $5.49 \times 10^{25}$

---

Q5.

A certain heat engine draws 500 cal/s from a water bath at 27 °C and transfers 400 cal/s to a reservoir at a lower temperature. The efficiency of this engine is:

- A) 20%  
B) 7.5%  
C) 55%  
D) 35%  
E) 10%

---

Q6.

A particle with a charge of  $5 \times 10^{-6}$  C and a mass of  $2.0 \times 10^{-2}$  kg moves with a constant speed of 7.0 m/s in a circular orbit around a stationary particle with a charge of  $-5 \times 10^{-6}$  C. The radius of the orbit is:

- A) 0.23 m  
B) 0.16 m  
C) 0.52 m  
D) 0.65 m  
E) 0.84 m

---

Q7.

A charged oil drop with a mass of  $2.00 \times 10^{-4}$  kg is held in equilibrium in air by a downward electric field of 300 N/C. The charge on the drop is:


- A)  $-6.53 \times 10^{-6}$  C  
B)  $-1.50 \times 10^{-6}$  C  
C)  $+6.53 \times 10^{-6}$  C  
D)  $+1.50 \times 10^{-6}$  C  
E)  $+3.57 \times 10^{-6}$  C

---

Q8.

Two identical large insulating parallel plates carry positive charge of equal magnitude that is distributed uniformly over their inner surfaces as shown in **Figure 2**. Rank the points 1 through 5 according to the magnitude of the electric field at these points, **least to greatest**.

Fig#



- A) (2 and 3) tie, then (1 and 4 and 5) tie
- B) 1, 2, 3, 4, 5
- C) 5, 4, 3, 2, 1
- D) (1 and 4 and 5) tie, then (2 and 3) tie
- E) (2 and 3) tie, then (1 and 4) tie, then 5

---

Q9.

An isolated solid metal sphere of radius  $R$  carries a charge of  $3.0 \text{ nC}$ . How much charge remains in the sphere of radius  $R$ , when it is connected to another uncharged metallic sphere of radius  $2R$  with a thin metallic wire? (Assume no charge remains on the wire and the spheres are far away from each other)

- A)  $1.0 \text{ nC}$
- B)  $2.0 \text{ nC}$
- C)  $6.0 \text{ nC}$
- D)  $1.5 \text{ nC}$
- E)  $3.0 \text{ nC}$

---

Q10.

Copper contains  $8.4 \times 10^{28}$  free electrons per cubic meter. A copper wire of radius  $5.0 \times 10^{-4} \text{ m}$  carries a current of  $1.0 \text{ A}$ . The electron drift speed is:

- A)  $9.5 \times 10^{-5} \text{ m/s}$
- B)  $1.0 \times 10^{-5} \text{ m/s}$
- C)  $6.5 \times 10^{-5} \text{ m/s}$
- D)  $3.0 \times 10^{-5} \text{ m/s}$
- E)  $5.0 \times 10^{-5} \text{ m/s}$

---

Q11.

A particle with a charge  $q_1 = 5.5 \times 10^{-8} \text{ C}$  is fixed at the origin. Another particle with a charge  $q_2 = -2.3 \times 10^{-8} \text{ C}$  is moved from position  $x = 3.5 \text{ cm}$  on the  $x$  axis to position  $y = 4.3 \text{ cm}$  on the  $y$  axis. Find the amount of work required to move the charge.

- A)  $+6.0 \times 10^{-5} \text{ J}$
- B)  $+3.1 \times 10^{-5} \text{ J}$
- C)  $-6.0 \times 10^{-5} \text{ J}$

- D)  $-3.1 \times 10^{-5} \text{ J}$
- E)  $+2.7 \times 10^{-5} \text{ J}$

Q12.


A parallel-plate capacitor has a plate area of  $0.30 \text{ m}^2$  and a plate separation of  $0.10 \text{ mm}$ . If the charge on each plate has a magnitude of  $5.0 \times 10^{-6} \text{ C}$ , what is the energy density in its electric field?

- A)  $16 \text{ J/m}^3$
- B)  $35 \text{ J/m}^3$
- C)  $78 \text{ J/m}^3$
- D)  $21 \text{ J/m}^3$
- E)  $54 \text{ J/m}^3$

Q13.

In **Figure 3** a battery with an emf  $\epsilon_1 = 12 \text{ V}$  and an internal resistance of  $r_1 = 1.0 \Omega$  is used to charge a battery with an emf  $\epsilon_2 = 7.0 \text{ V}$  and an internal resistance of  $r_2 = 1.0 \Omega$ . The current in the circuit is:

Fig#



- A) 2.5 A
- B) 1.0 A
- C) 3.0 A
- D) 4.1 A
- E) 7.5 A

Q14.


Two light bulbs, with power ratings  $40 \text{ W}$  and  $100 \text{ W}$ , are connected in series to a  $110 \text{ V}$  source. Then which of the following statements is **TRUE**?

- A) the current in the  $100 \text{ W}$  bulb is same as that in the  $40 \text{ W}$  bulb
- B) the current in the  $100 \text{ W}$  bulb is less than that in the  $40 \text{ W}$  bulb
- C) the voltage drop across the  $100 \text{ W}$  bulb is same as that in the  $40 \text{ W}$  bulb
- D) both bulbs have same energy dissipation rate
- E) the current in the  $100 \text{ W}$  bulb is greater than that in the  $40 \text{ W}$  bulb

Q15.

Find the equivalent resistance across points a and b in the circuit shown in **Figure 4**.

Fig#




- A) 2 Ω
- B) 4 Ω
- C) 1 Ω
- D) 6 Ω
- E) 3 Ω

Q16.

In the circuit diagram of **Figure 5**, if the current  $i = 0.5$  A, find the potential difference  $V_b - V_a$ .

Fig#



- A) 5.0 V
- B) 4.0 V
- C) 1.0 V
- D) 2.0 V
- E) 8.0 V

Q17.

A segment of a circuit diagram is shown in **Figure 6**. At a particular instant, if  $R = 2.0$  kΩ,  $C = 4.0$  mF,  $\epsilon = 8.0$  V,  $Q = 20$  mC, and  $i = 3.0$  mA, what is the potential difference  $V_a - V_b$ ?

Fig#




- A)  $-9.0 \text{ V}$
- B)  $+7.0 \text{ V}$
- C)  $-7.0 \text{ V}$
- D)  $+5.0 \text{ V}$
- E)  $+9.0 \text{ V}$

Q18.

An electron is travelling with constant velocity  $\vec{v}$  in a region of uniform electric field  $\vec{E}$  and the uniform magnetic field  $\vec{B}$ , as shown in **Figure 7**. Find the direction of the magnetic field  $\vec{B}$ .

Fig#



- A) the negative  $z$  direction (into the page)
- B) the negative  $y$  direction
- C) the positive  $y$  direction
- D) the positive  $z$  direction (out of the page)
- E) the negative  $x$  direction

Q19.


An electron has a velocity of  $6.0 \times 10^6 \text{ m/s}$  in the positive  $x$  direction at a point where the magnetic field has components  $B_x = 3.0 \text{ T}$ ,  $B_y = 1.5 \text{ T}$ , and  $B_z = 2.0 \text{ T}$ . What is the magnitude of the acceleration of the electron at this point?

- A)  $2.6 \times 10^{18} \text{ m/s}^2$
- B)  $3.4 \times 10^{18} \text{ m/s}^2$
- C)  $6.0 \times 10^{18} \text{ m/s}^2$
- D)  $1.2 \times 10^{18} \text{ m/s}^2$
- E)  $5.8 \times 10^{18} \text{ m/s}^2$

Q20.

**Figure 8** shows a loop of wire carrying a current  $i = 2.0$  A is in the shape of a right triangle with two equal sides, each 15 cm long. A uniform magnetic field  $B = 0.7$  T is in the plane of the triangle and is perpendicular to the hypotenuse. The resultant magnetic force on the two equal sides has a magnitude of:

Fig#



- A) 0.30 N
- B) 0.21 N
- C) 0.12 N
- D) 0.45 N
- E) 0.57 N

Q21.

A loop of current-carrying wire has a magnetic dipole moment of  $5.0 \times 10^{-4}$  A·m<sup>2</sup>. The dipole moment is initially aligned with a 0.50 T magnetic field. To rotate the loop so that its dipole moment becomes perpendicular to the field, you must do work of:

- A)  $+2.5 \times 10^{-4}$  J
- B) zero
- C)  $-2.5 \times 10^{-4}$  J
- D)  $+1.0 \times 10^{-3}$  J
- E)  $-1.0 \times 10^{-3}$  J

Q22.


A long wire of radius  $R = 4.5$  cm carries a uniform current throughout its cross-section. If the magnetic field inside the wire at 3.0 cm from the center is equal to three times the magnetic field at a distance  $r$  from the center, where  $r > R$ , calculate the distance  $r$ .

- A) 20 cm
- B) 35 cm
- C) 13 cm
- D) 46 cm
- E) 54 cm

Q23.

Two long vertical wires pierce (penetrate) the horizontal plane of the paper at the vertices of an equilateral triangle, each carrying 2.0 A current, one out of the paper and the other into the paper, as shown in **Figure 9**. The magnetic field at point P has a magnitude of:

Fig#



- A)  $1.0 \times 10^{-5}$  T
- B)  $8.2 \times 10^{-5}$  T
- C)  $1.7 \times 10^{-5}$  T
- D)  $5.5 \times 10^{-5}$  T
- E)  $2.9 \times 10^{-5}$  T

Q24.


Two long parallel wires X and Y are separated by 4.0 cm and carry currents 20 A and 30 A, respectively, along the same direction. Determine the magnitude of the magnetic force on a 2.0 m length of wire Y.

- A)  $6.0 \times 10^{-3}$  N
- B)  $4.0 \times 10^{-3}$  N
- C)  $2.0 \times 10^{-3}$  N
- D)  $3.0 \times 10^{-3}$  N
- E)  $7.0 \times 10^{-3}$  N

Q25.

In **Figure 10**, part of a long insulated wire carrying current  $i = 5.0$  A is bent into a circular section of radius  $R = 0.1$  m. What is the magnetic field at the center C of the circular section?

Fig#




- A)  $4.1 \times 10^{-5}$  T out of the page
- B)  $4.1 \times 10^{-5}$  T into the page
- C)  $5.5 \times 10^{-5}$  T into the page

- D)  $5.5 \times 10^{-5}$  T out of the page  
E)  $3.7 \times 10^{-5}$  T out of the page

Q26.

**Figure 11** shows the cross-sectional view of three wires carrying identical currents  $i$  and four Amperian loops (a through d) encircling them. Rank the loops according to the magnitude of  $\oint \vec{B} \cdot d\vec{s}$  along each, **greatest first**.

Fig#



- A) d, (a and c) tie, then b  
B) a, b, c and d  
C) d, (a and b) tie, then c  
D) c, (a and b) tie, d  
E) b, a, d, c

Q27.


A square loop (length along one side = 20 cm) rotates in a constant magnetic field which has a magnitude of 2.0 T. At an instant when the angle between the magnetic field and the normal to the plane of the loop is equal to  $20^\circ$  and increasing at a rate of 0.18 rad/s, what is the magnitude of the induced emf in the loop?

- A) 4.9 mV  
B) 1.3 mV  
C) 3.5 mV  
D) 2.1 mV  
E) 5.2 mV

Q28.

A long straight wire is in the plane of two circular conducting loops. The straight wire carries a constant current  $I$  in the direction shown in **Figure 12**. The circular loop 1 is moved to the right while the loop 2 is moved to the left with the same speed,  $v$ . The direction of the induced current in the circular loops 1 and 2 are respectively:

Fig#



- A) counter-clockwise , clockwise
- B) counter-clockwise , counter-clockwise
- C) clockwise , clockwise
- D) clockwise , counter-clockwise
- E) no direction because induced current is zero

Q29.


A long solenoid ( $n = 1500$  turns/m) has a cross-sectional area of  $0.40 \text{ m}^2$  and a current given by  $I = (4.0 + 3.0t^2) \text{ A}$ , where  $t$  is in seconds. A flat circular coil ( $N = 300$  turns) with a cross-sectional area of  $0.15 \text{ m}^2$  is inside and coaxial with the solenoid. What is the magnitude of the emf induced in the coil at  $t = 2.0 \text{ s}$ ?

- A) 1.0 V
- B) 2.0 V
- C) 1.5 V
- D) 2.5 V
- E) 3.5 V

Q30.

In the arrangement shown in **Figure 13**, a conducting bar of negligible resistance slides along horizontal, parallel, and frictionless conducting rails connected as shown to a  $2.0 \Omega$  resistor. A uniform  $1.5 \text{ T}$  magnetic field is perpendicular to the plane of the paper. If  $L = 60 \text{ cm}$ , at what rate is thermal energy being generated in the resistor at the instant the speed of the bar  $v = 4.2 \text{ m/s}$ ?

Fig#



- A) 7.1 W
- B) 2.6 W
- C) 5.0 W
- D) 1.2 W
- E) 3.6 W